

## **Comparative study of brain processes related to space-induced and clinical oculomotor disturbances**

### **Results:**

The studies included 3 groups of subjects: control group (C - 15 subjects), schizophrenic patients (S - 35 subjects), volunteers, exposed to 6-day dry immersion (DI - 12 subjects). All participants had right hand and right eye preference. Subjects performed visually-guided saccades and antisaccades. EEG was recorded from 19 standard sites. Mean amplitude of averaged potentials was assessed at 600 ms before the stimulus and at 600 to 100 ms before saccade onset.

The saccade characteristics did not differ in groups C and DI. Patients exhibited in the antisaccade task delays in the performance of correct saccades ( $p < 0,05$ ) and larger number of directional errors ( $p < 0,001$ ).

In control subjects the target stimulus was preceded by a vertex predominantly bilateral slow negative potential shift; left- and right-side saccades were preceded by PSN, that was larger at the midline and left sites, with parietal maximum before visually-guided saccades, and with frontal and parietal maximums before antisaccades. The PSN amplitude declined significantly in the DI and Sch groups as compared with the C group. However, in immersed subjects the foci of negativity shifted to the right hemisphere so that the PSN amplitude decreased sharply in the left and increased in the right hemisphere, with significant decrease in Fz, F3, Pz and P3. Significant decline of the PSN amplitude was the most characteristic in schizophrenic patients in the sagittal frontal and central regions (Fz and Cz).

The PSN shifts observed after exposure to simulated weightlessness and in schizophrenic patients could be caused by the alterations of sensory inputs' activities in weightlessness and fine frontal structural disorder of the brain in case of schizophrenia

### **Published work:**

Anna V. Kirenskaya, Vadim V. Myamlin, Vladimir Y. Novototsky-Vlasov, Mikhail V. Pletnikov, Inesa B. Kozlovskaya. (2011) The contingent negative variation laterality and dynamics in antisaccade task in normal and unmedicated schizophrenic subjects. *Span J Psychol.* 2011 Nov;14(2):869-83.

A.V.Kirenskaya, V.Yu.Novototsky-Vlasov, V.V.Myamlin, I.M.Ushakova. Characteristics of visually-guided saccades and presaccadic cortical negativity in unmedicated schizophrenic patients. 6<sup>th</sup> IBRO World Congress of Neuroscience. Prague, Czech Republic 2003, p. 354, ISBN 80-239-0887-1.

A.V.Kirenskaya, V.V.Myamlin, V.Yu.Novototsky-Vlasov, N.R.Gallyamova, I.M.Ushakova. The study of saccades' and presaccadic slow negative potentials' characteristics in schizophrenic patients. *Rossiyskiy psikhiatricheskiy zhurnal*, 2003, №6, pp. 23-30 (in Russian).

A.V.Kirenskaya, V.Yu.Novototsky-Vlasov, V.V.Myamlin, N.R.Gallyamova, I.M.Ushakova, I.B.Kozlovskaya. Slow presaccadic EEG-potentials in the antisaccade task in healthy subjects and in schizophrenic patients. *Rossiyskiy fiziologicheskiy zhurnal im. I.M.Sechenova*, 2004, V.90, № 8, p. 18 (in Russian).

V.Yu.Novototsky-Vlasov, M.V.Slavutskaya, A.V.Kirenskaya, V.V.Myamlin, N.R.Gallyamova. Slow brain potentials preceding saccades to visual stimuli in schizophrenic patients. *Rossiyskiy fiziologicheskiy zhurnal im. I.M.Sechenova*, 2004, V.90, № 8, p. 28 (in Russian).

E.S.Tomilovskaya, N.R.Gallyamova, V.Yu.Novototsky-Vlasov, A.V.Kirenskaya The influence of simulated micro gravitation on EEG characteristics, recorded during voluntary saccadic eye movements. *Rossiyskiy fiziologicheskiy zhurnal im. I.M.Sechenova*, 2004, V.90, № 8, p. 419 (in Russian).

I.B.Kozlovskaya, A.V. Kirenskaya, V.Yu.Novototsky-Vlasov, V.V.Myamlin, E.S.Tomilovskaya. Comparative study of the clinical (schizophrenia) and space-induced disturbances of event-related brain processes preceding voluntary saccades. 5° Simpósio da Fundação BIAL “Behind and beyond the brain”, Porto, 2004.

Tomilovskaya, E., Kirenskaya, A., Novototsky-Vlasov, V. Y. , & Kozlovskaya, I. (2004). Event-related EEG changes preceding saccadic eye movements before and after dry immersion. *Journal of Gravitational Physiology*, 11(2), 333-34.

E.S. Tomilovskaya, A.V. Kirenskaya, V.Yu. Novototsky-Vlasov, I.B. Kozlovskaya. EEG spectral power dynamic changes related to saccadic eye movements before and after dry immersion. – In: 25<sup>th</sup> Annual International Gravitational Physiology Meeting, 2004, Russian Academy of Sciences, Moscow, Russia

Kirenskaya, A., Kamenskov, M., Myamlin, V., Novototsky-Vlasov, V. Y., & Tkachenko, A. (2013). The antisaccade task performance deficit and specific CNV abnormalities in patients with stereotyped paraphilia and schizophrenia. *Journal of Forensic Sciences*, 58(5), 1219–1226. doi: 10.1111/1556-4029.12241

Slavutskaya, M., Kirenskaya, A., Novototsky-Vlasov, V. Y., Shulgovsky, V. & Kozlovskaya, I. (2005). Slow cortical potentials preceding visually guided saccades in schizophrenics. *Human physiology*, 31(5), 545-553. doi: 10.1007/s10747-005-0095-z

## **Researchers' Contacts:**

Inesa B. Kozlovskaya  
5 Dolgorukovskaya st., ap.287, 127006  
Moscow, Russia

Telephone: +7-095-251-1491  
E-mail: inesa@ibmp.ru